arXiv:1501.01356v3 [math.GR] 9 May 2015

Permutation-like Matrix Groups with a Maximal Cycle of Power of Odd Prime Length

Guodong Deng, Yun Fan

School of Mathematics and Statistics
Central China Normal University, Wuhan 430079, China

Abstract

If every element of a matrix group is similar to a permutation matrix, then it is called a permutation-like matrix group. References [3] and [4] showed that, if a permutation-like matrix group contains a maximal cycle of length equal to a prime or a square of a prime and the maximal cycle generates a normal subgroup, then it is similar to a permutation matrix group. In this paper, we prove that if a permutation-like matrix group contains a maximal cycle of length equal to any power of any odd prime and the maximal cycle generates a normal subgroup, then it is similar to a permutation matrix group.

Key words: permutation-like matrix group, permutation matrix group, maximal cycle.

MSC2010: 15A18, 15A30, 20H20.

1 Introduction

Let $\mathbb{C}$ be the complex filed. Let $\mathrm{GL}_d(\mathbb{C})$ be the complex general linear group of dimension d, i.e. the multiplicative group consisting of invertible complex $d \times d$ matrices. Any subgroup $\mathcal{G} \le \mathrm{GL}_d(\mathbb{C})$ is called a matrix group of dimension d. For a matrix group $\mathcal{G}$, if there exists a $P \in \mathrm{GL}_d(\mathbb{C})$ such that $P^{-1}AP$ for all $A \in \mathcal{G}$ are permutation matrices, then $\mathcal{G}$ is said to be *similar (or conjugate) to a permutation matrix group*, or said to be a *permutation matrix group* for short. If for every element A of $\mathcal{G}$ there exists an $S \in \mathrm{GL}_d(\mathbb{C})$ such that $S^{-1}AS$ is a permutation matrix, then $\mathcal{G}$ is called a *permutation-like matrix group*.

Cigler [2, 3] showed that a permutation-like matrix group is not a permutation matrix group in general. A $d \times d$ matrix is called a *maximal cycle* if it is similar to a permutation matrix corresponding to the cycle permutation of length d. Cigler conjectured that:

Email address: yfan@mail.ccnu.edu.cn (Yun Fan).

Conjecture: If a permutation-like matrix group contains a maximal cycle, then it is a permutation matrix group.

Cigler [2, 3] proved that, if a permutation-like matrix group $\mathcal{G}$ of dimension prime p contains a maximal cycle which generates a normal cyclic subgroup, then $\mathcal{G}$ is a permutation matrix group. In [4], we further proved that this result still holds if we replace the dimension prime p by dimension p^2.

In this paper we prove:

Theorem 1.1. *Let $\mathcal{G}$ be a permutation-like matrix group of dimension p^n where p is any odd prime and n is any positive integer. If $\mathcal{G}$ contains a maximal cycle which generates a normal cyclic subgroup, then $\mathcal{G}$ is a permutation matrix group.*

Necessary preparations for the proof of the theorem are made in Section 2. For fundamentals of the group theory, please refer to [1, 5]. In Section 3 we treat a special case of the theorem where $\mathcal{G}$ is a p-group, i.e. its order is a power of p. The proof of Theorem 1.1 is presented in Section 4.

2 Preparation

We begin with few preliminaries on relations between a cyclic group $\langle C\rangle$ of order d generated by C and the residue ring $\mathbb{Z}_d$ of the integer ring $\mathbb{Z}$ modulo the positive integer d. We have a bijection which maps $k \in \mathbb{Z}_d$ to $C^k \in \langle C\rangle$. By $\mathbb{Z}_d^*$ we denote the multiplicative group consisting of the reduced residue classes in $\mathbb{Z}_d$. Obviously, $\langle C^k\rangle = \langle C\rangle$ if and only if $k \in \mathbb{Z}_d^*$; at that case we say that C^k is a *generator* of the cyclic group $\langle C\rangle$. Any automorphism α of $\langle C\rangle$ is corresponding to exactly one $r \in \mathbb{Z}_d^*$ such that $\alpha(C^k) = C^{rk}$ for any $C^k \in \langle C\rangle$, By μ_r we denote the permutation of $\mathbb{Z}_d$ mapping $k \in \mathbb{Z}_d$ to $\mu_r(k) = rk$. The automorphism group of $\langle C\rangle$ is isomorphic to $\mathbb{Z}_d^*$ by mapping α to μ_r as above. If $\mathcal{G}$ is a finite group which contains $\langle C\rangle$ as a normal subgroup, then $\mathcal{G}$ is homomorphic to a subgroup of $\mathbb{Z}_d^*$ with kernel consisting of the elements which centralize $\langle C\rangle$.

For an element A of a group $\mathcal{G}$, we denote the order of A by $\mathrm{ord}_{\mathcal{G}}(A)$, or $\mathrm{ord}(A)$ for short if the group is known from context. Let p be a prime. A finite group $\mathcal{G}$ is called a *p-group* if its order $|\mathcal{G}|$ is a power of p. The next lemma about a cyclic p-group is obvious. As mentioned above, it is also a lemma about $\mathbb{Z}_{p^n}$. Note that $|\mathbb{Z}_{p^n}^*| = p^{n-1}(p-1)$.

Lemma 2.1. *Let $\mathcal{G} = \langle C\rangle$ be a cyclic group and $|\mathcal{G}| = p^n$.*

(i) For $0 \le a < n$ denote $\mathcal{G}^{p^a} = \{C^{p^a t} \mid 0 \le t < p^{n-a}\}$. Then $\mathcal{G}^{p^a} = \langle C^{p^a}\rangle$ is a cyclic subgroup of $\mathcal{G}$ of order p^{n-a} generated by C^{p^a}.

(ii) The map $\mathcal{G} \to \mathcal{G}^{p^a}$, $C^k \mapsto C^{p^a k}$, is an epimorphism with kernel $\mathcal{G}^{p^{n-a}}$. In particular, for any generator G of the cyclic group $\mathcal{G}^{p^a}$ there are exactly p^a generators of $\mathcal{G}$ which are mapped to G.

(iii) If p is an odd prime and α is an automorphism of $\mathcal{G}$, then $\mathrm{ord}(\alpha) = sp^a$ with $s \mid (p-1)$ and $0 \le a < n$, and there are integers u, v coprime to p such that $\mathrm{ord}_{\mathbb{Z}_{p^n}^*}(u) = s$ (hence $\mathrm{ord}_{\mathbb{Z}_p^*}(u) = s$) and $\alpha(C) = C^{u+vp^{n-a}}$. □

For a subgroup $\mathcal{H}$ of a group $\mathcal{G}$, if any element which centralizes $\mathcal{H}$ is contained in $\mathcal{H}$, then $\mathcal{H}$ is said to be *self-centralized*.

Lemma 2.2. *Assume that $\mathcal{G}$ is a finite group containing a self-centralized normal cyclic p-subgroup $\langle C\rangle$ generated by an element C, where p is an odd prime. Then $\mathcal{G} = \langle A, C\rangle$ generated by two elements, and one of the following two holds.*

(i) If $\mathcal{G}$ is a p-group, then there is an $A' \in \mathcal{G}$ such that $\mathcal{G} = \langle A', C\rangle$ and $\langle A'\rangle \cap \langle C\rangle = 1$.

(ii) If $\mathcal{G}$ is not a p-group, then $\langle A\rangle \cap \langle C\rangle = 1$.

Proof. We sketch a proof, some of the arguments will be quoted later.

Assume that $|\langle C\rangle| = p^n$. Since $\langle C\rangle$ is self-centralized, the quotient group $\mathcal{G}/\langle C\rangle \cong \langle A\rangle/\langle A\rangle \cap \langle C\rangle$ is isomorphic to a subgroup of the multiplicative $\mathbb{Z}^*_{p^n}$. The group $\mathbb{Z}^*_{p^n}$ is a cyclic group of order $p^{n-1}(p-1)$. Thus, $|\mathcal{G}/\langle C\rangle| = sp^a$ with $0 \le a < n$ and $s \mid (p-1)$, and there is an $r \in \mathbb{Z}^*_{p^n}$ such that

$$A^{-1}CA = C^r \quad \text{and} \quad \operatorname{ord}_{\mathbb{Z}^*_{p^n}}(r) = |\mathcal{G}/\langle C\rangle| = sp^a. \tag{2.1}$$

For any integer k and positive integer j, it is a direct computation that

$$(AC^k)^j = (AC^k)\cdots(AC^k)(AC^k) = A^j C^{k(r^{j-1}+\cdots+r+1)}. \tag{2.2}$$

(i). Assume that $|\mathcal{G}/\langle C\rangle| = p^a$, $0 < a < n$. By Lemma 2.1(iii), we assume that $r = 1 + vp^{n-a}$ with $p \nmid v$. Denote $\langle C\rangle^A = \{C^t \mid A^{-1}C^tA = C^t\}$. Note that $A^{-1}C^tA = C^{tr}$. So $A^{-1}C^tA = C^t$ if and only if $t(r-1) \equiv 0 \pmod{p^n}$. Thus $\langle C\rangle^A = \langle C^{p^a}\rangle$. Since $\mathcal{G}/\langle C\rangle \cong \langle A\rangle/\langle A\rangle \cap \langle C\rangle$, $A^{p^a} \in \langle C\rangle$ hence $A^{p^a} \in \langle C\rangle^A = \langle C^{p^a}\rangle$. So, we can find an integer k such that $A^{p^a}C^{kp^a} = 1$. Note that

$$r^{p^a-1} + \cdots + r + 1 = \frac{r^{p^a}-1}{r-1} = \frac{(1+vp^{n-a})^{p^a}-1}{vp^{n-a}} \equiv p^a \pmod{p^n}. \tag{2.3}$$

Let $A' = AC^k$. By Eqns (2.2) and (2.3), we obtain

$$A'^{p^a} = A^{p^a}C^{k(r^{p^a-1}+\cdots+r+1)} = A^{p^a}C^{kp^a} = 1.$$

Thus, $\mathcal{G} = \langle A', C\rangle$ and $\langle A'\rangle \cap \langle C\rangle = 1$.

(ii). Assume that $|\mathcal{G}/\langle C\rangle| = sp^a$, $0 \le a < n$ and $s \mid (p-1)$. Since $\mathcal{G}$ is not a p-group, we have $s > 1$. By Lemma 2.1 (iii), we can assume that $\operatorname{ord}_{\mathbb{Z}^*_p}(r) = s$. Then $p \nmid (r-1)$ (otherwise $s = 1$). As we have seen, $A^{-1}C^tA = C^t$ if and only if $t(r-1) \equiv 0 \pmod{p^n}$. At the present case, $t(r-1) \equiv 0 \pmod{p^n}$ if and only if $t \equiv 0 \pmod{p^n}$. Thus $\langle C\rangle^A = 1$. As argued above, $A^{sp^a} \in \langle C\rangle^A$. So $A^{sp^a} = 1$. That is, $\langle A\rangle \cap \langle C\rangle = 1$. □

We turn to preliminaries on matrices.

A diagonal blocked matrix $\begin{pmatrix} B_1 & & \\ & \ddots & \\ & & B_m \end{pmatrix}$ is denoted by $B_1 \oplus \cdots \oplus B_m$ for short. The identity matrix of dimension d is denoted by $I_{d\times d}$, or I for short if the dimension is known from context.

We denote the characteristic polynomial of a complex matrix A by $\operatorname{char}_A(x)$.

Lemma 2.3 ([4, Lemma 2.1])**.** *The following two are equivalent to each other:*

(i) A *is similar to a permutation matrix;*

(ii) A *is diagonalizable and* $\mathrm{char}_A(x) = \prod_i (x^{\ell_i} - 1)$.

If it is the case, then each factor $x^{\ell_i} - 1$ *of* $\mathrm{char}_A(x)$ *corresponds to exactly one* ℓ_i*-cycle of the cycle decomposition of the permutation of the permutation matrix.* $\square$

By $\Phi_n(x)$ we denote the cyclotomic polynomial of degree n, i.e. $\Phi_n(x) = \prod_\omega (x-\omega)$ with ω running over the primitive n-th roots of unity. Since $x^n - 1 = \prod_{k|n} \Phi_k(x)$, the following is an immediate consequence of the above lemma.

Corollary. *Let* A *be a matrix similar to a permutation matrix, and* m, n *be positive integers. If* $\Phi_n(x)^m \,\big|\, \mathrm{char}_A(x)$*, then* $\Phi_k(x)^m \,\big|\, \mathrm{char}_A(x)$ *for any* $k \mid n$. $\square$

The next lemma is a combination of [4, Eqns (2.1), (2.2), (2.3) and (2.4)].

Lemma 2.4. *Let* $C \in \mathrm{GL}_d(\mathbb{C})$ *be a maximal cycle of dimension* d*, and* λ *be a primitive* d*-th root of unity. Then the following hold.*

(i) $\{\lambda^j \mid j \in \mathbb{Z}_d\} = \{\lambda^0 = 1, \lambda, \cdots, \lambda^{d-1}\}$ *is the spectrum (i.e. the set of eigenvalues) of* C.

(ii) The eigen-subspace of every eigenvalue λ^j *of* C*, denoted by* $\mathrm{E}(\lambda^j)$*, has dimension* 1*, and* $\mathbb{C}^d = \bigoplus_{j=0}^{d-1} \mathrm{E}(\lambda^j)$.

(iii) If $e_j \in \mathrm{E}(\lambda^j)$ *for* $j = 0, 1, \cdots, d-1$ *are all nonzero, then* $(e_0, e_1, \cdots, e_{d-1})$ *is a basis of* $\mathbb{C}^d$ *and* $Ce_j = \lambda^j e_j$, $j = 0, 1, \cdots, d-1$.

(iv) Let $f = \alpha_0 e_0 + \alpha_1 e_1 + \cdots + \alpha_{d-1} e_{d-1}$ *where* $e_0, \cdots, e_{d-1}$ *are as above and all* $\alpha_j \in \mathbb{C}$*. Then* $(f,\ Cf,\ \cdots,\ C^{d-1}f)$ *is a basis of* $\mathbb{C}^d$ *if and only if* $\alpha_j \neq 0$ *for all* $j = 0, 1, \cdots, d-1$*. If it is the case, with respect to that basis,* C *is a cycle permutation matrix of dimension* d. $\square$

Lemma 2.5 ([4, Lemma 2.2])**.** *Let* C *and* λ *be as above in Lemma 2.4. Let* $A \in \mathrm{GL}_d(\mathbb{C})$ *such that* $A^{-1}CA = C^r$ *for an* $r \in \mathbb{Z}_d^*$*. Further assume that* $\mathbb{Z}_d$ *is partitioned into* μ_r*-orbits as follows:*

$$\Gamma_0 = \{0\},\ \Gamma_1 = \{j_1, rj_1, \cdots, r^{d_1-1}j_1\},\ \cdots,\ \Gamma_m = \{j_m, rj_m, \cdots, r^{d_m-1}j_m\}$$

i.e. $r^{d_k} j_k \equiv j_k \pmod d$ *but* $r^{d_k-1} j_k \not\equiv j_k \pmod d$*. For* $k = 0, \cdots, m$*, take nonzero* $e_{j_k} \in \mathrm{E}(\lambda^{j_k})$*, set* $\mathcal{E}_k = \{e_{j_k}, Ae_{j_k}, \cdots, A^{d_k-1}e_{j_k}\}$ *and* $\mathcal{E} = \bigcup_{k=0}^m \mathcal{E}_k$*. Then the following hold.*

(i) $A \cdot \mathrm{E}(\lambda^j) = \mathrm{E}(\lambda^{rj})$, $j = 0, \cdots, d-1$*, where* $A \cdot \mathrm{E}(\lambda^j) = \{Av_j | v_j \in \mathrm{E}(\lambda^j)\}$.

(ii) $\mathcal{E}_k$ *is a basis of* $V_k := \bigoplus_{h=0}^{d_j-1} \mathrm{E}(\lambda^{r^h j_k})$*, and* A *restricted to* V_k *has the matrix*

$$A|_{\mathcal{E}_k} = \begin{pmatrix} 0 & \cdots & 0 & \omega_k \\ 1 & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 \end{pmatrix}_{d_k \times d_k}, \tag{2.4}$$

where ω_k is an $(\mathrm{ord}(A)/d_k)$*-th root of unity,*

(iii) $\mathbb{C}^d = V_1 \oplus \cdots \oplus V_m$*, the union* $\mathcal{E} = \mathcal{E}_1 \cup \cdots \cup \mathcal{E}_m$ *is a basis of* $\mathbb{C}^d$ *and, with respect to the basis* $\mathcal{E}$*, the matrix of* A *is*

$$A|_{\mathcal{E}} = A|_{\mathcal{E}_1} \oplus \cdots \oplus A|_{\mathcal{E}_m}. \qquad \square$$

Proposition 2.6 ([4, Proposition 2.3])**.** *Let the notations be as in Lemma 2.5. If the matrix* $A|_{\mathcal{E}}$ *is a permutation matrix (i.e. all* $\omega_k = 1$*), then the matrix group* $\langle A, C\rangle$ *generated by* A *and* C *is a permutation matrix group.* $\square$

Note that "the matrix $A|_{\mathcal{E}}$ is a permutation matrix" means that mapping $e \in \mathcal{E}$ to Ae is a permutation of the set $\mathcal{E}$.

Lemma 2.7 ([3, Proposition 4.2])**.** *If* $\mathcal{G}$ *is a permutation-like matrix group and* $C \in \mathcal{G}$ *is a maximal cycle, then* $\langle C\rangle$ *is self-centralized in* $\mathcal{G}$*.*

We sketch the proof for convenience. For $A \in \mathcal{G}$ which centralizes C, we can assume that C is a permutation matrix of dimension d corresponding to the d-cycle permutation and $A = \sum_{i=0}^{d-1} \alpha_i C^i$. Since AC^{d-k} is similar to a permutation matrix, its trace $d\alpha_k$ is a non-negative integer. Hence α_k for $k = 0, 1, \cdots, d-1$ are non-negative rationals. By Lemma 2.3, all the eigenvalues of A are roots of unity, in particular, $\sum_{i=0}^{d-1} \alpha_i = 1 = \big|\sum_{i=0}^{d-1} \alpha_i \lambda^i\big|$. If there are at least two of the coefficients non-zero, say $\alpha_k \neq 0 \neq \alpha_h$ for $0 \le k \neq h < d$, then $\big|\alpha_k\lambda^k + \alpha_h\lambda^h\big| < \alpha_k + \alpha_h$ because $\lambda^k \neq \lambda^h$; hence

$$1 = \Big|\sum_{i=0}^{d-1} \alpha_i\lambda^i\Big| \le \big|\alpha_k\lambda^k + \alpha_h\lambda^h\big| + \sum_{i\neq k,h} \big|\alpha_i\lambda^i\big| < \sum_{i=1}^{d-1} \alpha_i = 1,$$

which is impossible. Thus there is exactly one of the coefficients, say α_k, which is non-zero; then $\alpha_k = 1$ and $A = C^k$.

3 The case of p-groups

In the following p is always an odd prime and n is a positive integer. In this section we consider a special case of Theorem 1.1, where the permutation-like matrix group is a p-group. This is a key step for the proof of the theorem.

Let $C \in \mathrm{GL}_{p^n}(\mathbb{C})$ be a maximal cycle. We have a disjoint union $\mathbb{Z}_{p^n} = p\mathbb{Z}_{p^n} \bigcup \mathbb{Z}_{p^n}^*$, where

$$p\mathbb{Z}_{p^n} = \{pk \pmod{p^n} \mid k \in \mathbb{Z}_{p^n}\} = \{0, p, \cdots, p(p^{n-1}-1)\}.$$

By Lemma 2.4, we have two subspaces of $\mathbb{C}^{p^n}$, denoted by V^p and V^*, as follows:

$$V^p = \bigoplus_{j\in p\mathbb{Z}_{p^n}} \mathrm{E}(\lambda^j) = \bigoplus_{e\in\mathcal{E}^p} \mathbb{C}e \quad \text{and} \quad V^* = \bigoplus_{j\in \mathbb{Z}_{p^n}^*} \mathrm{E}(\lambda^j) = \bigoplus_{e\in\mathcal{E}^*} \mathbb{C}e \tag{3.1}$$

where $\mathcal{E}^p$ and $\mathcal{E}^*$ denote the basis of V^p and V^* respectively as in Lemma 2.5. Then $\mathbb{C}^{p^n} = V^p \oplus V^*$.

For any $A \in \mathrm{GL}_{p^n}(\mathbb{C})$ which normalizes $\langle C \rangle$, there is exactly one $r \in \mathbb{Z}_{p^n}^*$ such that $A^{-1}CA = C^r$. Both $p\mathbb{Z}_{p^n}$ and $\mathbb{Z}_{p^n}^*$ are μ_r-invariant. So, for any integer k with $0 < k < p^n$ both V^p and V^* are AC^k-invariant subspaces of $\mathbb{C}^{p^n}$. By $AC^k|_{V^p}$ and $AC^k|_{V^*}$ we denote the linear transformations of AC^k restricted to V^p and V^* respectively. Correspondingly, $AC^k|_{\mathcal{E}^p}$ and $AC^k|_{\mathcal{E}^*}$ are matrices of $AC^k|_{V^p}$ and $AC^k|_{V^*}$ respectively.

Lemma 3.1. *Let $A, C \in \mathrm{GL}_{p^n}(\mathbb{C})$. Assume that C is a maximal cycle and $A^{-1}CA = C^r$ for an $r \in \mathbb{Z}_{p^n}^*$ with $\mathrm{ord}_{\mathbb{Z}_{p^n}^*}(r) = p^a$ where $0 \le a < n$. Let V^p and V^* be as in Eqn* (3.1)*. If $A^{p^a} = I$, then for*

$$\mathrm{char}_{(AC^k)|_{V^*}}(x) = \begin{cases} \Phi_{p^{n-\nu_p(k)}}(x)^{p^{\nu_p(k)}}, & 0 \le \nu_p(k) < n-a; \\ (x^{p^a}-1)^{p^{n-a-1}(p-1)}, & \nu_p(k) \ge n-a, \text{ or } k = 0; \end{cases}$$

where $\nu_p(k)$ denotes the p-adic valuation of k, i.e. $p^{\nu_p(k)}$ is the largest power of p which divides k.

Proof. Assume that $\mathbb{Z}_{p^n}^*$ is partitioned into μ_r-orbits $\Gamma_1, \cdots, \Gamma_h$. Every orbit Γ_i has length p^a, and the number $h = p^{n-a-1}(p-1)$. We can assume that

$$\Gamma_1 = \{j_1, rj_1, \cdots, r^{p^a-1}j_1\}, \ \cdots, \ \Gamma_h = \{j_h, rj_h, \cdots, r^{p^a-1}j_h\}.$$

For the basis $\mathcal{E}^*$ of V^*, by Lemma 2.5 we have $\mathcal{E}^* = \mathcal{E}_1^* \cup \cdots \cup \mathcal{E}_h^*$ with

$$\mathcal{E}_i^* = \{e_{j_i}, \ Ae_{j_i}, \ \cdots, \ A^{r^{p^a-1}}e_{j_i}\}, \qquad i = 1, \cdots, h. \tag{3.2}$$

Denote the restricted matrices to $\mathcal{E}_i^*$ by $A_i = A|_{\mathcal{E}_i^*}$ and $C_i = C|_{\mathcal{E}_i^*}$ for $i = 1, \cdots, h$. Since $A_i^{p^a} = I$, we have

$$A_i = \begin{pmatrix} 0 & \cdots & 0 & 1 \\ 1 & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 \end{pmatrix}_{p^a \times p^a}, \ C_i = \begin{pmatrix} \lambda^{j_i} & & & \\ & \lambda^{rj_i} & & \\ & & \ddots & \\ & & & \lambda^{r^{p^a-1}j_i} \end{pmatrix}_{p^a \times p^a}. \tag{3.3}$$

Thus $AC^k|_{\mathcal{E}^*} = \bigoplus_{i=1}^h A_iC_i^k$, and

$$\mathrm{char}_{A_iC_i^k}(x) = x^{p^a} - \lambda^{\sum_{j \in \Gamma_i} jk} = x^{p^a} - \lambda^{j_ik(1+r+\cdots+r^{p^a-1})}.$$

The conclusion is obviously true if $k = 0$. So we further assume that $k \neq 0$. By Lemma 2.1 (iii), we can take an integer v which is coprime to p such that $r = 1 + vp^{n-a}$. Then it is easy to check that $\nu_p(1 + r + \cdots + r^{p^a-1}) = a$, see

Eqn (2.3). So we can write $1 + r + \cdots + r^{p^a - 1} = a'p^a$ and $k = k'p^{\nu_p(k)}$ with $p \nmid a'$ and $p \nmid k'$. Then

$$\mathrm{char}_{A_i C_i^k}(x) = x^{p^a} - \lambda^{a'k'j_i p^{\nu_p(k)+a}}, \qquad i = 1, \cdots, h.$$

By Lemma 2.1 (ii), the collection of $\lambda^{a'k'j_i p^{\nu_p(k)+a}}$ for $i = 1, \cdots, h$ is just the collection of all primitive $p^{n-a-\nu_p(k)}$-th roots of unity, each of which appears with multiplicity

$$\frac{h}{p^{n-a-\nu_p(k)-1}(p-1)} = p^{\nu_p(k)}.$$

If $n - a - \nu_p(k) > 0$, then the collection of roots of $\mathrm{char}_{AC^k|_{V^*}}(x)$ is just the collection of all primitive $p^{n-\nu_p(k)}$-th roots of unity, each of which has multiplicity $p^{\nu_p(k)}$; hence

$$\mathrm{char}_{AC^k|_{V^*}}(x) = \Phi_{p^{n-\nu_p(k)}}(x)^{p^{\nu_p(k)}}.$$

Otherwise, $n - a - \nu_p(k) \le 0$, i.e. $\lambda^{a'k'j_i p^{\nu_p(k)+a}} = 1$, hence

$$\mathrm{char}_{AC^k|_{V^*}}(x) = (x^{p^a} - 1)^h = (x^{p^a} - 1)^{p^{n-a-1}(p-1)}. \qquad \square$$

Corollary 3.2. *Let notation be as in Lemma 3.1. If the matrix group* $\langle A, C\rangle$ *is permutation-like and* $A^{p^a} = I$*, then* $\langle A|_{V^p}, C|_{V^p}\rangle$ *is a permutation-like matrix group of dimension* p^{n-1}*.*

Proof. Let ℓ, k be any non-zero integers. Let $A' = A^\ell$, $a' = a - \nu_p(\ell)$ and $r' = r^\ell$. Then $A'^{-1}CA' = C^{r'}$ and $\mathrm{ord}(A') = \mathrm{ord}_{\mathbb{Z}_{p^n}^*}(r') = p^{a'}$. by Lemma 3.1,

$$\mathrm{char}_{(A^\ell C^k)|_{V^*}}(x) = \begin{cases} \Phi_{p^{n-\nu_p(k)}}(x)^{p^{\nu_p(k)}}, & \nu_p(k) < n - a'; \\ (x^{p^{a'}} - 1)^{p^{n-a'-1}(p-1)}, & \nu_p(k) \ge n - a'. \end{cases}$$

Since $A^\ell C^k = A'C^k$ is similar to a permutation matrix, by Lemma 2.3 and its corollary,

$$\mathrm{char}_{A^\ell C^k|_{V^p}}(x) = \begin{cases} (x^{p^{n-\nu_p(k)-1}} - 1)^{p^{\nu_p(k)}}, & \nu_p(k) < n - a'; \\ \prod_i (x^{p^i} - 1)^{j_i}, & \nu_p(k) \ge n - a'. \end{cases}$$

By Lemma 2.3 again, the matrix group $\langle A|_{V^p}, C|_{V^p}\rangle$ is a permutation-like matrix group of dimension p^{n-1}. $\square$

Lemma 3.3. *Let* $A, C \in \mathrm{GL}_{p^n}(\mathbb{C})$*. Assume that* C *is a maximal cycle and* $A^{-1}CA = C^r$ *for an* $r \in \mathbb{Z}_{p^n}^*$ *with* $\mathrm{ord}_{\mathbb{Z}_{p^n}^*}(r) = p$*. If* $\langle A, C\rangle$ *is a permutation-like matrix group and* $A^p = I$*, then* $A|_{V^p} = I$*.*

Proof. Note that $\langle A, C\rangle = \{A^\ell C^k \mid 0 \leq \ell < p,\ 0 \leq k < p^n\}$. For any $0 < \ell < p$ and $0 < k < p^n$, by Lemma 3.1 (note that $a = 1$ at present case),

$$\mathrm{char}_{A^\ell C^k|_{V^*}}(x) = \begin{cases} \Phi_{p^{n-\nu_p(k)}}(x)^{p^{\nu_p(k)}}, & \nu_p(k) < n-1; \\ (x^p-1)^{p^{n-2}(p-1)}, & \nu_p(k) \geq n-1. \end{cases}$$

Since $A^\ell C^k$ is similar to a permutation matrix, by Lemma 2.3,

$$\mathrm{char}_{A^\ell C^k|_{V^p}}(x) = \begin{cases} \left(x^{p^{n-\nu_p(k)-1}} - 1\right)^{p^{\nu_p(k)}}, & \nu_p(k) < n-1; \\ (x-1)^{p^{n-1}-pj}(x^p-1)^j, & \nu_p(k) \geq n-1. \end{cases} \tag{3.4}$$

Thus $\left\langle A|_{\mathcal{E}^p}, C|_{\mathcal{E}^p}\right\rangle$ is a permutation-like matrix group of dimension p^{n-1}.

Obviously, $C|_{\mathcal{E}^p} = \bigoplus_{j \in p\mathbb{Z}_{p^n}} \lambda^j$ is a maximal cycle of dimension p^{n-1}. Since $r = 1 + vp^{n-1}$ for some integer v coprime to p (see Lemma 2.1(iii)), for any $pt \in p\mathbb{Z}_{p^n}$ we have $\mu_r(pt) \equiv pt \pmod{p^n}$. So, $A|_{\mathcal{E}^p}$ is a diagonal matrix, hence $A|_{\mathcal{E}^p}$ commutes with $C|_{\mathcal{E}^p}$. By Lemma 2.7, $A|_{\mathcal{E}^p} \in \left\langle C|_{\mathcal{E}^p}\right\rangle$. But $(A|_{\mathcal{E}^p})^p = I$ and $\mathrm{ord}(C|_{\mathcal{E}^p}) = p^{n-1}$. Thus $A|_{\mathcal{E}^p} \in \left\langle C_{\mathcal{E}^p}^{p^{n-2}}\right\rangle$, and we can assume that $A|_{\mathcal{E}^p} = (C|_{\mathcal{E}^p})^{bp^{n-2}}$ with $0 \leq b < p$. Suppose that $b > 0$, then $\nu_p(-bp^{n-2}) = n-2$, and by Eqn (3.4),

$$\mathrm{char}_{I_{p^{n-1}\times p^{n-1}}}(x) = \mathrm{char}_{(A|_{\mathcal{E}^p})(C|_{\mathcal{E}^p})^{-bp^{n-2}}}(x) = (x^p-1)^{p^{n-2}}.$$

However, $\mathrm{char}_{I_{p^{n-1}\times p^{n-1}}}(x) = (x-1)^{p^{n-1}}$. This is a contradiction. Thus $b = 0$ and $A|_{\mathcal{E}^p} = I_{p^{n-1}\times p^{n-1}}$. □

Proposition 3.4. *Let $A, C \in \mathrm{GL}_{p^n}(\mathbb{C})$. Assume that C is a maximal cycle and $A^{-1}CA = C^r$ for an $r \in \mathbb{Z}_{p^n}^*$ with $\mathrm{ord}_{\mathbb{Z}_{p^n}^*}(r) = p^a$ where $0 \leq a < n$. If $\mathcal{G} = \langle A, C\rangle$ is a permutation-like matrix group and $A^{p^a} = I$, then $A|_{\mathcal{E}}$ is a permutation matrix.*

Proof. If $n = 1$, then $a = 0$ and $A = I$, the proposition holds trivially.

Assume $n > 1$. From Eqn (3.3) we have seen that $A|_{V^*}$ is a permutation matrix. By Corollary 3.2, $\langle A|_{V^p}, C|_{V^p}\rangle$ is a permutation-like matrix group of dimension p^{n-1}. Note that $\mathrm{ord}_{\mathcal{G}}(A^{p^{a-1}}) = \mathrm{ord}_{\mathbb{Z}_{p^n}^*}(r^{p^{a-1}}) = p$. By Lemma 3.3, $(A|_{V^p})^{p^{a-1}} = I$. And, $\mathrm{ord}_{\mathbb{Z}_{p^{n-1}}^*}(r) = p^{a-1}$. By induction on n, $A|_{\mathcal{E}^p}$ is a permutation matrix. Hence $A|_{\mathcal{E}} = A|_{\mathcal{E}^p} \oplus A|_{\mathcal{E}^*}$ is a permutation matrix. □

Corollary 3.5. *Let $A, C \in \mathrm{GL}_{p^n}(\mathbb{C})$. Assume that C is a maximal cycle and A normalizes $\langle C\rangle$. If $\mathcal{G} = \langle A, C\rangle$ is a permutation-like matrix p-group, then $\mathcal{G}$ is a permutation matrix group.*

Proof. By Lemma 2.7, $\langle C\rangle$ is self-centralized in $\mathcal{G}$. By Lemma 2.2(i), we have an $A' \in \mathcal{G}$ such that $\mathcal{G} = \langle A', C\rangle$, $A'^{-1}CA' = C^r$ for an $r \in \mathbb{Z}_{p^n}^*$ with $\mathrm{ord}_{\mathbb{Z}_{p^n}^*}(r) = p^a$ and $A'^{p^a} = I$. Thus, by Proposition 3.4 and Proposition 2.6, $\mathcal{G}$ is a permutation group. □

4 Proof of Theorem 1.1

Let $\mathcal{G} \le \mathrm{G}_{p^n}(\mathbb{C})$ be a permutation-like matrix group of dimension p^n where p is an odd prime. Let $C \in \mathcal{G}$ be a maximal cycle such that $\langle C\rangle$ is a normal subgroup of $\mathcal{G}$. By Lemma 2.7, $\langle C\rangle$ is self-centralized in $\mathcal{G}$. By Lemma 2.2 and Eqn (2.1), $\mathcal{G} = \langle A, C\rangle$, $A^{-1}CA = C^r$ where $r \in \mathbb{Z}_{p^n}^*$ with

$$\mathrm{ord}_{\mathbb{Z}_{p^n}^*}(r) = |\mathcal{G}/\langle C\rangle| = sp^a, \qquad s \,|\, (p-1), \quad 0 \le a < n.$$

Let $\mathcal{E}$ be the basis of $\mathbb{C}^{p^n}$ as in Lemma 2.5.

Case 1: $s = 1$. So $\mathcal{G}$ is a p-group, and Theorem 1.1 holds by Corollary 3.5.

Case 2: $a = 0$ and $s > 1$. By Lemma 2.1 (iii), we can assume that $\mathrm{ord}_{\mathbb{Z}_p^*}(r) = s$. Then for any $0 < \ell < s$, $r^\ell \not\equiv 1 \pmod p$, i.e. $p \nmid (r^\ell - 1)$. Thus $r^\ell k \equiv k \pmod{p^n}$ if and only if $k \equiv 0 \pmod{p^n}$. So every μ_r-orbit on $\mathbb{Z}_{p^n}$ has length s except for the orbit $\{0\}$. By Lemma 2.2 (ii), $A^s = I$. By Lemma 2.5 (ii) and (iii),

$$A|_{\mathcal{E}} = \omega_0 \oplus \begin{pmatrix} 0 & \cdots & 0 & 1 \\ 1 & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 \end{pmatrix}_{s\times s} \oplus \cdots \oplus \begin{pmatrix} 0 & \cdots & 0 & 1 \\ 1 & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 \end{pmatrix}_{s\times s}.$$

And the characteristic polynomial

$$\mathrm{char}_A(x) = (x - \omega_0)(x^s - 1)^{(p^n-1)/s}.$$

By Lemma 2.3, $\omega_0 = 1$. So $A|_{\mathcal{E}}$ is a permutation matrix. By Proposition 2.6, we obtain that $\mathcal{G}$ is a permutation matrix group.

Case 3: $a > 0$ and $s > 1$. Then $\mathcal{G}$ is not a p-group. By Lemma 2.2(ii) we further have that $A^{sp^a} = I$. Since s and p^a are coprime each other, we have integers t, m such that $st + p^a m = 1$. Let $A' = A^{st}$ and $A'' = A^{p^a m}$. Then $A = A^{st+p^a m} = A'A''$, $A'^{p^a} = I$ and $A''^s = I$. By Proposition 3.4, $A'|_{\mathcal{E}}$ is a permutation matrix. From the above argument of Case 2, $A''|_{\mathcal{E}}$ is also a permutation matrix. Thus $A|_{\mathcal{E}} = A'|_{\mathcal{E}} \cdot A''|_{\mathcal{E}}$ is a permutation matrix. By Proposition 2.6, $\mathcal{G}$ is a permutation matrix group. $\square$

Acknowledgments

The authors are supported by NSFC through the grant number 11271005. Thanks are given to the reviewers for the helpful suggestions.

References

[1] J. L. Alperin, R. B. Bell, Groups and Representations, GTM 162, Springer-Verlag, New York, 1997.

[2] G. Cigler, Groups of matrices with prescribed spectrum, Doctoral dissertation, 2005, http://matknjiz.si/doktotati/2005/10921-83.pdf

[3] G. Cigler, Permutation-like matrix groups, Linear Algebra and its Applications 422(2007), 486-505.

[4] Guodong Deng, Yun Fan, Permutation-like matrix group with a maximal cycle of prime square length, Linear Algebra and its Applications 450(2014), 44-55.

[5] D.J.S. Robinson, A Course in Groups Theory, GTM 80, Springer-Verlag, New York, 1980.